arXiv:1106.5276v2 [math.GR] 25 Jul 2011

FINE ASYMPTOTIC GEOMETRY IN THE HEISENBERG GROUP

MOON DUCHIN AND CHRISTOPHER MOONEY

Abstract. For every finite generating set on the integer Heisenberg group $H(\mathbb{Z})$, Pansu showed that the word metric has the large-scale structure of a Carnot-Carathéodory Finsler metric on the real Heisenberg group $H(\mathbb{R})$. We study the properties of those limit metrics and obtain results about the geometry of word metrics that reflect the dependence on generators.

For example we will study the probability that a group element has all of its geodesic spellings sublinearly close together, relative to word length. In free abelian groups of rank at least two, that probability is 0; in unbounded hyperbolic groups, the probability is 1. In $H(\mathbb{Z})$ it is a rational number strictly between 0 and 1 that depends on the generating set; with respect to the standard generators, the probability is precisely 19/31.

1. Introduction

In this paper we will focus on the 3-dimensional integer Heisenberg group $H(\mathbb{Z})$, a uniform lattice in the real Heisenberg group $H(\mathbb{R})$, with the word metric coming from a finite generating set. We will study large-scale geometric properties that are sensitive to the choice of generators, such as the shape and stability of long geodesics, and asymptotic density of classes of elements.

We will take an approach based on work of Pansu and Breuillard [9, 2] exploiting the fact that all word metrics on $H(\mathbb{Z})$ have the large-scale structure of Carnot-Carathéodory Finsler metrics on $H(\mathbb{R})$; that is, the Lie group with its cc metric is the asymptotic cone of the lattice with the chosen word metric. However, we will study large-scale geometric properties that are lost when one passes to the asymptotic cone. (Note that studying the Heisenberg group up to quasi-isometry is even coarser, identifying all word metrics as well as all left-invariant metrics on $H(\mathbb{R})$ with one another.) Thus we develop technology for working with these word metrics that uses the homogeneous dilation at finite scale together with combinatorial arguments, instead of the usual tools of coarse geometry. We think of this set of questions and techniques as belonging to "fine asymptotic geometry."

The geometry of nilpotent groups is closely tied to geometric minimax problems, so Section 2 is devoted to the solution of relevant isoperimetric problems in normed planes. In Section 3, we apply those findings, giving a very explicit description of the limit metric associated to an arbitrary word metric. The main result of this section is Theorem 10, which is summarized here:

Structure Theorem. *For any finite symmetric generating set S of $H(\mathbb{Z})$, the limiting* cc *metric admits a complete description of its geodesics, classified into no more than*

Date: July 27, 2011.

The first author was partially supported by NSF DMS-0906086. The second author was partially supported by NSF RTG-0602191.

$|S|^2 - 2|S|$ *combinatorial types of* regular geodesics *and no more than* $|S|$ *types of* unstable geodesics. *The unit sphere* $\mathcal{S}$ *in the* CC *metric is a piecewise union of the graphs of quadratic polynomials over finitely many quadrilateral regions in the xy-plane.*

As a corollary of this description, we can analyze the uniqueness of geodesics in these limit metrics, describing exactly which points $\mathsf{x} \in H(\mathbb{R})$ are reached by more than one geodesic segment based at $\mathsf{0}$.

We use this structure theorem in Section 4 to understand the geodesics in Cayley graphs for $H(\mathbb{Z})$, showing that word geodesics may be very unruly, but they are tracked to within a controlled distance by much better-behaved CC geodesics: the Hausdorff distance is of lower-order growth than the length of the word. (See Lemma 18.)

Tracking Lemma. *Away from a certain unstable locus of endpoints, word geodesics are sublinearly tracked by* CC *geodesics.*

This opens the door to the study of geometric probability in the discrete Heisenberg groups, and we give an illustrative application in Theorem 22, quantifying the instability of word geodesics. Elements of a group may be said to be geodesically stable with respect to a generating set if, in the Cayley graph, the Hausdorff distance between the geodesic spellings of the word can only differ from each other sublinearly in the length of the word. When this is made precise, one can see that stable elements have zero density in free abelian groups, but full density in infinite hyperbolic groups. The situation is different in the Heisenberg group.

Instability Theorem. *The asymptotic density of geodesically stable elements of* $H(\mathbb{Z})$ *is a rational number strictly between* 0 *and* 1*, depending on the generating set.*

Then in Section 5, we apply the structure theorem to a generalized Gauss circle problem: Theorem 24 asserts that in every polygonal CC metric—a class that includes all limits of word metrics—we can count lattice points to first order in the annular shell between balls.

Counting Theorem. *For any polygonal* CC *metric, the number of lattice points between the spheres of radius* n *and* $n-1$ *is equal to* $4Vn^3 + O(n^2)$*, where* V *is the volume of the* CC *unit ball.*

Note that this is strictly better than what is logically implied by the best known estimate of the number of lattice points in the ball of radius n, which is $Vn^4 + O(n^3)$ by [12]. (In fact, the full theorem is more precise than what is stated above: we can count points in the annulus in any radial direction.)

Finally, we show in Section 6 that for $S = \mathsf{std}$, the cardinality of the discrete sphere S_n agrees with this annular lattice point count to first order. The counting measure on discrete spheres is thus shown to limit to a cone measure on the limit shape $\mathcal{S}$, for which we have a concrete and finite description. This enables spherical averaging—sharper than asymptotic density calculations, which are averages over balls—with respect to the standard generators.

1.1. **Nilpotent groups and the Heisenberg group.** We review a few facts about the sub-Riemannian geometry of the Heisenberg group; see [5] for a comprehensive reference.

First, we consider the 3-dimensional integer Heisenberg group $H(\mathbb{Z})$, the subgroup of matrices with integer entries in $H(\mathbb{R})$ (the real Heisenberg group). Let $\mathfrak{m}$ denote the horizontal subspace of the Lie algebra $\mathfrak{h}$ of $H(\mathbb{R})$; that is, the span of the tangent

vectors $X_0 = \left(\begin{smallmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{smallmatrix}\right)$ and $Y_0 = \left(\begin{smallmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{smallmatrix}\right)$. There are *horizontal planes* at every point in $H(\mathbb{R})$, generated by pushing around X_0 and Y_0 by left multiplication in $H(\mathbb{R})$ to produce left-invariant vector fields X and Y. We say that a curve in $H(\mathbb{R})$ is *admissible* if all of its tangent vectors lie in these horizontal planes; it is easily verified that the curve $\gamma = (\gamma_1, \gamma_2, \gamma_3)$ is admissible if and only if

$$\gamma_3' = \frac{1}{2}(\gamma_1\gamma_2' - \gamma_2\gamma_1').$$

We will use the *exponential coordinates* on H given by the following representation:

$$(x, y, z) \leftrightarrow \begin{pmatrix} 1 & x & z + \frac{1}{2}xy \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}.$$

These coordinates have the property that $(x, y, z)^n = (nx, ny, nz)$.

For integers x and y, define $\epsilon(x, y)$ to be $1/2$ if x and y are both odd, and 0 otherwise. In these coordinates, $H(\mathbb{Z})$ looks just like the standard lattice $\mathbb{Z}^3 \subset \mathbb{R}^3$ shifted by ϵ in the z direction.

We can regard $\mathfrak{m}$ as a copy of $\mathbb{R}^2$ and make use of the linear projection $\pi : H(\mathbb{R}) \to \mathfrak{m}$ given by $(x, y, z) \mapsto (x, y)$.

Lemma 1 (Standard fact from Heisenberg geometry)**.** *For any path $\gamma = (\gamma_1, \gamma_2)$ from $(0, 0)$ to (x, y), there is a unique admissible curve $\overline{\gamma} = (\gamma_1, \gamma_2, \gamma_3)$. The lifted curve connects the origin to the point (x, y, z), where z is equal to the signed Euclidean area of R, the region in the plane enclosed by γ and a straight chord from $(0, 0)$ to (x, y).*

Proof. To construct the lift, we just need to supply the coordinate function γ_3, which can be done by integrating in the defining formula for admissibility. Note that $x(t)y'(t) - y(t)x'(t)$ is identically zero along a linear parametrization of the straight chord between the endpoints, so we can concatenate that chord with the path in the plane to obtain the closed curve ∂R. We get

$$z = \int_{\partial R} \gamma_1\gamma_2' - \gamma_2\gamma_1' = \int_R dx \wedge dy,$$

which is the area of R. □

From now on the area of this region R will be called the *balayage area* associated to a curve γ.

Choose a centrally symmetric convex body $Q \subset \mathfrak{m}$ with boundary L and let $\|\cdot\|_L$ denote the corresponding norm; this is the norm having Q as its unit ball and L as its unit sphere. We will use the notation $\mathfrak{m}_L = (\mathfrak{m}, \|\cdot\|_L)$ for this normed plane. This induces a Carnot-Carathéodory Finsler metric (from now on, CC metric) on the Lie group: the distance between two points is the infimal length of an admissible path between them, measured by using the norm on the tangent vectors. (This is well defined for admissible curves because the tangent vectors lie in copies of $\mathfrak{m}$ pushed around by multiplication in $H(\mathbb{R})$, and it is a classical fact that this gives $H(\mathbb{R})$ the structure of a geodesic space.) Note that measuring the length of an admissible curve in $H(\mathbb{R})$ with respect to this CC metric is equivalent to measuring the length of its projection to $\mathfrak{m}$ with respect to the norm $\|\cdot\|_L$. If L is a polygon, we will call the induced metric a *polygonal* CC *metric* on $H(\mathbb{R})$. (In arbitrary dimension, if L is a polytope, the associated norms are sometimes called *crystalline*.)

These CC metrics are equipped with a dilation $\delta_t(x,y,z) = (tx, ty, t^2 z)$ that scales lengths and distances by t, and areas in $\mathfrak{m}$ by t^2. For any set $E \subset H(R)$, let ΔE denote its full cone under dilation, and let $\hat{E}$ denote its dilation-cone to the origin:

$$\Delta E = \{\delta_t(\mathsf{x}) : \mathsf{x} \in E, t \geq 0\} \quad ; \qquad \hat{E} = \{\delta_t(\mathsf{x}) : \mathsf{x} \in E, 0 \leq t \leq 1\}.$$

As a consequence of the connection between height and balayage area, we have a criterion for geodesity in the CC metric: a curve γ in $\mathfrak{m}$ based at $(0,0)$ lifts to a geodesic in $H(\mathbb{R})$ iff its L-length is minimal among all curves with the same endpoints and enclosing the same area.

We will consider geometric probability, or *asymptotic density*, in the Heisenberg groups; for $U \subset H(\mathbb{Z})$ and $V \subset H(\mathbb{R})$, let $B_t = B_t(\mathsf{0})$ be the ball of radius t about the origin. Then

$$\mathrm{Prob}(U) := \lim_{n\to\infty} \frac{|B_n \cap U|}{|B_n|} \quad ; \qquad \mathrm{Prob}(V) := \lim_{r\to\infty} \frac{\mathrm{vol}(B_r \cap V)}{\mathrm{vol}(B_r)}.$$

Note that Lebesgue measure has the right invariance properties to coincide with Haar measure on $H(\mathbb{R})$ up to a scalar, so we will be able to talk about ratios of volumes unambiguously.

1.2. **Limit shapes, limit metrics, and limit measure.** We have the following extremely general theorem, first proven for nilpotent groups by Pansu, and extended to all periodic pseudometrics on simply connected solvable Lie groups of polynomial growth by Breuillard [2]. Here we just state it for the very special case of word metrics on $H(\mathbb{Z}) \leq H(\mathbb{R})$. We will always assume that generating sets are symmetric ($S = -S$).

Theorem 2 (Pansu [9])**.** *Consider a word metric on $H(\mathbb{Z})$ given by a finite generating set S. Let $\pi(S)$ be the linear projection of the generators S to the horizontal subspace $\mathfrak{m}$, let Q be the convex hull of $\pi(S)$ and L its boundary polygon. Then the limit*

$$\mathcal{S} = \lim_{n\to\infty} \delta_{\frac{1}{n}} S_n \ ,$$

exists (as a Gromov-Hausdorff limit, say) and is equal to the unit sphere in the CC metric induced by the norm $\|\cdot\|_L$ on $\mathfrak{m}$.

Equivalently,

$$\lim_{\mathsf{x}\to\infty} \frac{|\mathsf{x}|_S}{d_{\mathrm{cc}}(\mathsf{x},\mathsf{0})} = 1.$$

Theorem 3 (Krat [8])**.** *Furthermore, there exists a constant $K = K(S)$ such that*

$$d_{\mathrm{cc}}(\mathsf{x},\mathsf{0}) - K \ \leq \ |\mathsf{x}|_S \ \leq \ d_{\mathrm{cc}}(\mathsf{x},\mathsf{0}) + K$$

for all $\mathsf{x} \in H(\mathbb{Z})$.

The second result, bounded difference between the word metric and the CC metric, is stronger here in $H(\mathbb{Z})$ than for the general case, where one only has that the ratio goes to 1. In fact Breuillard showed in [2] that there exist 2-step nilpotent groups where bounded difference fails.

For the word metric $(H(\mathbb{Z}), S)$, we will call $\mathcal{S}$ and the induced CC metric the *limit shape* and *limit metric*, respectively. The volume of the unit ball in the CC metric will turn out to be an important number to attach to both the word metric and the CC metric itself; we will denote it by

$$V = V(S) = V(L) := \mathrm{vol}(\hat{\mathcal{S}}).$$

The Hausdorff dimension of $H(\mathbb{R})$ is equal to 4, despite the topological dimension of 3, as one can see clearly by considering the growth of a cube under the dilation. Thus the volume of the ball of radius r in the CC metric equals $V \cdot r^4$, and in $H(\mathbb{Z})$, we have $|B_n| = V \cdot n^4 + O(n^3)$.

In the previous paper [7], the case of $\mathbb{Z}^d$ was studied in detail, and there we also obtained a *limit measure* for a generating set S. In that case Q is the convex hull in $\mathbb{R}^d$ of the generators S themselves, and L is again its boundary. The limit shape here is given by $L = \lim_{n\to\infty} \frac{1}{n} S_n$. In [7] it was shown that counting measure on $\frac{1}{n}S_n$ converges to a measure on L called *cone measure*, which assigns to a measurable set $\sigma \subset L$ the measure $\mu_L(\sigma) = \frac{\mathrm{vol}(\hat{\sigma})}{\mathrm{vol}(Q)}$. (That is, it is the proportion of the area of Q that is subtended by σ, in this case by Euclidean dilation.) This allows us to reduce the problem of averaging over larger and larger S_n with respect to counting measure to the finite problem of averaging over L against cone measure.

Below, in Theorem 25, we will obtain the analogous result for $H(\mathbb{Z})$ with respect to the standard generators, which enables us to calculate spherical averages; conjecturally, cone measure is the limit measure for all finite generating sets.

1.3. **Relationship to prior literature on $H(\mathbb{Z})$.** Pansu's seminal paper from 1983 [9], based on his dissertation, shows that the ratio of limit metric to word metric goes to one for all virtually nilpotent groups with finite generating sets, as discussed above, and concludes that these word metrics on $H(\mathbb{Z})$ satisfy $|B_n| = \alpha \cdot n^d + o(n^d)$. Breuillard's preprint from 2007 [2] extends this result (that the growth has a well-defined leading coefficient) to a much larger class of groups, and computes the coefficient. Breuillard also gives extremely concrete geometric constructions in that paper, especially in the Appendix, where some explicit computations are shown for $(H(\mathbb{Z}), \mathsf{std})$ that inspired the current work.

M. Shapiro wrote a study of $H(\mathbb{Z})$ with respect to its standard generators in 1989 [10], giving a description of S_n, then computing an exact formula for $|S_n|$ as a polynomial function of n. This remarkable formula has not only a well-defined leading coefficient, but in fact all of the coefficients are computed and only the constant term oscillates (in fact, between twelve different values). He uses this study to establish that $H(\mathbb{Z})$ has infinitely many cone types, and is almost convex in the sense of Cannon.

Stoll 1998 [12] refines Pansu's asymptotics by showing that for 2-step nilpotent groups with finite generating sets, $|B_n| = \alpha \cdot n^d + O(n^{d-1})$, and this error term is "often" sharp—compare our counting theorem (Thm 24). In a separate paper, Stoll had showed that 2-step nilpotent groups with infinite cyclic derived subgroup have a finite generating set for which the growth series is rational; on the other hand, higher Heisenberg groups (of dimension 5 and up) have a finite generating set for which the growth series is transcendental [11]. These results are recovered by Breuillard by geometric methods.

Blachère 2003 [1] computes an exact word length formula with respect to the standard generators, using it to show "almost-connectedness" of discrete spheres S_n. We note that the calculations of Shapiro and Blachère for S_n with respect to $S = \mathsf{std}$ are somewhat combinatorial, and that they are essentially equivalent to the more geometric description of S_n given below in §6 (which also give directional information).

Finally, in work based on her dissertation, Dani 2007 [6] has some striking results on asymptotic density: for any virtually nilpotent group, she gives an exact formula

for the density of finite-order elements. Further, she shows that the values of these densities range over $\mathbb{Q} \cap [0,1]$ as the groups vary.

Acknowledgments. Many thanks to Ralf Spatzier, Alex Eskin, and Emmanuel Breuillard for useful conversations on numerous aspects of the problems considered here.

2. Isoperimetry and isoarea in normed planes

2.1. **The classical problem.** As seen above, understanding CC geodesics in $H(\mathbb{R})$ amounts to the problem of finding paths in $\mathfrak{m}$ of minimal L-length given their endpoints and balayage area. For closed loops, this is just the classical isoperimetric problem in the normed space $\mathfrak{m}_L$, which was elegantly solved by Busemann in the 1940s as follows.

Let the *polar dual* Q^* of a convex, centrally symmetric polygon Q be defined by

$$Q^* = \{\mathsf{x} \in \mathfrak{m} : \ \mathsf{x} \cdot \mathsf{y} \le 1 \quad \forall \mathsf{y} \in Q\}.$$

Then if u and v are successive vertices of Q, Q^* has a vertex a corresponding to the edge between them which is the solution to $\mathsf{a} \cdot \mathsf{u} = \mathsf{a} \cdot \mathsf{v} = 1$. (That is, if the side is called σ and the line through the origin perpendicular to σ is called ℓ, then the vertex dual to σ is a vector pointing in the direction of ℓ and whose length is the reciprocal of the length of the projection of u or v to ℓ.) In particular, if Q is a polygon with $2N$ sides, then Q^* is also a $2N$-gon.

Theorem 4 (Busemann [4])**.** *Consider the norm on $\mathbb{R}^2$ induced by a convex centrally symmetric body Q. Then the maximal ratio of enclosed area to perimeter is uniquely achieved (up to scale) by the* isoperimetrix *I, which is defined to be the boundary of the rotate by $\pi/2$ of the polar dual Q^* of Q.*

If Q is a polygon, then the sides of I are parallel to the vertex directions of Q. Here is the calculation: suppose a and b are successive vertices of Q^*, so they are dual to successive edges of Q, sharing a vertex v. Then a side of I is obtained by rotating $a - b$ by $\pi/2$, so that side of I is perpendicular to $a - b$. But since $a \cdot v = 1$ and $b \cdot v = 1$ (by construction of a and b), we have $(a - b) \cdot v = 0$, which shows that the side of I is parallel to v.

Most of the rest of this section is devoted to solving modifications of this problem that are relevant for the nilpotent geometry. We will introduce classes of curves called *trace paths* and *beelines*, then show that these provide solutions to the Dido problem and the dual isoarea problem in $\mathfrak{m}_L$.

2.2. **Trace paths.** Let I_1 be a scaled copy of I, scaled to have perimeter one in the L-norm. Fix a parametrized 1-periodic curve $\iota : \mathbb{R} \to \mathbb{R}^2$ so that the image of ι is I_1, the parametrization is by arclength relative to the norm, and ι traverses I_1 counterclockwise.

For any choice of parameters $t \in [0,1]$ and $T \in (t, t+1]$, let

$$\mathsf{trace}_{t,T}(s) = \frac{\iota\big(sT + t - st\big) - \iota(t)}{T - t}, \qquad s \in [0,1].$$

We call these *trace paths*: they are parametrized paths of length one in the norm, whose shape follows along the perimeter of $\frac{1}{T-t} I_1$ (a scaled copy of the isoperimetrix). These trace paths start at the origin and therefore must terminate inside the unit ball in the norm (namely Q). See Figure 1 for a classification of trace paths for two different choices of Q.

Definition 5. *For a given normed plane $\mathfrak{m}_L$, let the* balayage function $A : Q \to \mathbb{R}$ *denote the maximal balayage area among all paths of length* 1 *from* $(0,0)$ *to* (x,y).

Lemma 6 (Trace paths solve the Dido problem)**.** *Given $(x,y) \in Q$, a path γ of L-length* 1 *connecting* $(0,0)$ *and* (x,y) *has maximal balayage area among all such paths if and only if it is a trace path: $\gamma = \mathsf{trace}_{t,T}$ where $\mathsf{trace}_{t,T}(1) = (x,y)$.*

Proof. Consider a particular trace path with $\mathsf{trace}_{t,T}(1) = (x,y)$. Let α be the straight chord from $(0,0)$ to (x,y) and let $\lambda = \frac{1}{T-t}$. Let A_{tr} denote the area enclosed by $\mathsf{trace}_{t,T}$ and α; we want to show that this is maximal. Note that the trace path can be completed to λI_1 (a scaled copy of I) by extending the domain up to $s = \lambda$. Thus α can be realized as a chord of λI_1 that cuts it into two pieces, and A_{tr} is the area of one of those two pieces. But now suppose there were a different path γ in $\mathbb{R}^2$, of length one in the L-norm, connecting $(0,0)$ to (x,y) while enclosing maximal area $A(x,y) \geq A_{\mathsf{tr}}$ with the chord α. Maximality of area implies convexity, so in particular the path γ always stays on one side of the line through α. But then γ can also be concatenated with $\mathsf{trace}_{t,T}([1,\lambda])$, producing a figure which must be a copy of λI_1, by uniqueness of the isoperimetrix. Thus $A_{\mathsf{tr}} = A(x,y)$, showing that trace paths enclose maximal area among all paths of length one, as desired. Indeed, this also means that the image of γ is uniquely determined, which forces it to be a trace path, though it is possible that $\gamma = \mathsf{trace}_{t,T} = \mathsf{trace}_{t',T'}$ for another choice of parameters with $t' = t + k$, $T' = T + k$. □

From here on we will assume that Q is a $2N$-gon, in which case I_1 will also be a $2N$-gon and ι will be piecewise affine: $\iota(s) = s\mathsf{a} + \mathsf{b}$ for fixed vectors a, b when s is in the subinterval of values corresponding to a side of I. Let ι be chosen so that $\iota(0)$ is a vertex in a copy of I_1 and let $\sigma_1, \sigma_2, \ldots, \sigma_{2N}$ be the sides listed counterclockwise. Let ℓ_i be the sidelength of σ_i for all i, so that $\sum \ell_i = 1$.

Define

$$Q_{ij} := \left\{ \frac{\iota(T) - \iota(t)}{T - t} : \quad \iota(t) \in \sigma_i, \quad \iota(T) \in \sigma_j \right\}.$$

This is the subset of Q consisting of endpoints $\mathsf{trace}_{t,T}(1)$ of those trace paths whose shape on I_1 begins on the ith side and ends on the jth side.

We will say that two continuous, piecewise linear paths have the *same shape* if there exist vector directions $\mathsf{v}_1, \ldots, \mathsf{v}_k$ such that each path follows $a_1\mathsf{v}_1, a_2\mathsf{v}_2, \ldots, a_k\mathsf{v}_k$ in the same order, with $a_1, a_k \geq 0$, $a_2, \ldots, a_{k-1} > 0$, and $\mathsf{v}_i \neq \mathsf{v}_{i+1}$ for all i.

Theorem 7 (Combinatorics of trace paths)**.** *If Q is a polygon with $2N$ sides, then there are $4N^2 - 4N$ quadrilaterals Q_{ij}, covering Q and overlapping only on their boundaries, such that the trace paths have "constant combinatorics" on each piece: that is, the trace paths ending at any two points in Q_{ij} have the same shape.*

Proof. We will show that the quadrilaterals Q_{ii} and $Q_{i,i+1}$ are degenerate (have no area in $\mathfrak{m}$). Discarding those, there remain $(2N)(2N-2)$ nondegenerate quadrilaterals, which will be seen to have the properties described in the statement above.

It will be helpful to note right away that within a given Q_{ij}, each level set in t or T is a straight line. This is because ι is affine on each edge. Thus, within the quadrilaterals and for fixed T, the function trace is of the form $(t\mathsf{a} + \mathsf{b})/(c - t)$, so its derivative is $\frac{1}{(c-t)^2}[c\mathsf{a} + \mathsf{b}]$, which has constant direction. This shows that t-level sets (and, similarly, T-level sets) are a family of straight segments sweeping out each quadrilateral, though not in general a parallel family. Let us also consider level sets of trace at $T - t = k$. In



FIGURE 1. The combinatorics of trace paths for a square and a hexagon. We can count 8 and 24 nondegenerate quadrilaterals, respectively, with constant combinatorics on each.

this case we have

$$\mathsf{trace}_{t,t+k}(1) = \frac{\iota(t+k) - \iota(t)}{k},$$

which produces a parallel family of straight segments within Q_{ij} indexed by k.

We now consider paths starting and ending on the ith side. This quadrilateral Q_{ii} has two components: either the whole trace path is contained in the ith side, or the trace path traverses all the sides of I before returning. Recall that the side σ_i is parallel to a vertex direction from L, say for vertex v_i. If the trace path is totally contained in σ_i then the image $\mathsf{trace}_{t,T}(1)$ is the vertex v_i itself, because it points in that direction and has length one in the L norm. In the other case, we point in the $-\mathsf{v}_i$ direction. The magnitude of the vectors in this component of Q_{ii} range from 0 (achieved when $T = 1 + t$, so the path closes up) to $\ell_i/(1-\ell_i)$ (achieved when t is at the end of σ_i and T has wrapped around to the beginning of σ_i). Thus each Q_{ii} is a vertex of L together with a line segment in the interior of Q.

Now we will show that $Q_{i,i+1}$ is the edge of L with endpoints v_i, v_{i+1}. This follows simply from noticing that each of those trace paths is a two-sided path following direction v_i for time a and then direction v_{i+1} for time $1-a$, so that it terminates at the point $a\mathsf{v}_i + (1-a)\mathsf{v}_{i+1}$. Notice that these particular trace paths are geodesic in $\mathbb{R}^2$ with respect to the L-norm; and indeed since all trace paths have L-length one, they are L-norm-geodesics if and only if they terminate on L itself (the sphere of radius one).

For the remaining quadrilaterals, there are four extreme points: we may start at either endpoint of σ_i and terminate at either endpoint of σ_j. Indeed, there is a quadrilateral traced out by holding the start point or end point on I_1 fixed at one extreme while moving the other from extreme to extreme, then alternating which is fixed and which is moving, making a circuit of length four. (Its four subpaths are level sets for t or T, so they are straight lines.)

Finally, observe that $\mathsf{trace}_{t,t+k}(1)$ is a convex, centrally symmetric polygon for every fixed $k \in (0,1]$. This polygon varies continuously in k, is identically zero when $k = 1$ and equals L itself for k sufficiently small. We can conclude that all points in Q are hit by trace paths. □

2.3. **Beelines.** Let us call a path in $\mathbb{R}^2$ a *beeline* if it has L-length one and connects the origin to a point on L. That is, beelines are L-geodesic segments emanating from the origin. One can check that a given path ending interior to a side σ of L is a beeline by verifying that all of its tangent vectors point towards σ. The name reflects their consistent progress away from the origin, as opposed to trace paths which typically make turns in several directions that seem inefficient in the L-norm.

Lemma 8 (Existence of beelines)**.** *Every point $(x,y) \in L$ is reached by a unique trace path $\mathsf{trace}_{t,T}$, and this path encloses a nonnegative area $A = A(x,y)$. There are beelines to (x,y) enclosing every signed area in the range $[-A, A]$.*

Proof. If (x,y) is in the edge of L with extreme points v_i and v_{i+1}, then the trace path to (x,y) follows two directions, first going distance a in direction v_i and then distance $(1-a)$ in direction v_{i+1}. Now vary s from 0 to a and consider the beeline that goes s in direction v_i followed by $(1-a)$ in direction v_{i+1} and finally $(a-s)$ in direction v_i. This is clearly still an L-geodesic, since it has length one and terminates on L. But the area it encloses with the straight chord between its endpoints varies continuously from $-A$ when $s = 0$ to A when $s = a$. □

In general, the full set of beelines to a side with extreme points v_i and v_{i+1} consists of parametrizations by arclength of arbitrary piecewise differentiable paths whose tangent vectors lie in the interval of directions between those two extremes.

Lemma 9 (Isoarea problem)**.** *A curve in Q of L-length 1 has minimal length among all curves with the same endpoints and balayage area if and only if it is a trace path or a beeline.*

Proof. The reverse implication is easy: we have already shown that trace paths uniquely maximize area to their endpoints. Since beelines are L-norm geodesics, there is no shorter path between their endpoints.

Now for the forward implication: Let β be a curve in Q of L-length 1 which has minimal length among all curves with the same endpoints and balayage area. If β ends on L, then it is a beeline by definition. Otherwise say β ends at $(x, y) \in Q^\circ$ and has balayage area A'. We know that there is a trace path $\gamma = \mathsf{trace}_{t,T}$ with $\mathsf{trace}_{t,T}(1) = (x, y)$ enclosing area $A(x, y) \geq A'$. If $A(x, y) = A'$, then β is a trace path by Lemma 6. If not, then we can modify γ to be an improvement on β: we know that the region enclosed by γ is convex, so we can find a chord of this body starting and ending on γ so that the modified curve γ', which follows γ except for a shortcut along this chord, has length < 1 and encloses area A'. This contradicts the minimality of the length of β. □

3. Structure of polygonal cc metrics

3.1. **Unit sphere and cc geodesics.** Returning to the Heisenberg geometry, we find that we have identified the geodesics, and can thus map the shape of spheres.

Recall that Pansu's theorem tells us that the word metric $(H(\mathbb{Z}), S)$ is asymptotic to the cc norm on $H(\mathbb{R})$ induced by L, the boundary of the convex hull of $\pi(S)$. Note that the number of sides of L is at most $|S|$, but could be smaller if S has several elements with the same projection to $\mathfrak{m}$, or has elements which do not project to vertices of L. If L is a $2N$-gon, then there are $4N^2 - 4N$ quadrilateral regions Q_{ij} classifying the shapes of trace paths, while the $2N$ sides of L classify the beelines.

Theorem 10 (Structure Theorem)**.** *Consider the cc metric induced on $H(\mathbb{R})$ by a polygonal norm $\|\cdot\|_L$ on $\mathfrak{m}$. The balayage function $A(x, y)$ is a continuous function over Q whose restriction to each Q_{ij} is a quadratic polynomial in x, y.*

Let $\mathbf{z}(x, y)$ be the multivalued function given by

$$\mathbf{z} = \begin{cases} \{\pm A\}, & (x, y) \in Q^\circ \\ [-A, A], & (x, y) \in L. \end{cases}$$

Then the graph of $\mathbf{z}$ is precisely the unit sphere $\mathcal{S}$ in the cc metric.

Equivalently: the full set of cc geodesics of length one based at the origin is identical to the set of admissible lifts of trace paths and beelines (and their reflections in the origin).

Proof. For points $(x, y, z) \in H(\mathbb{R})$ with $z \geq 0$ and $(x, y) \in Q^\circ$, saying that an admissible path α from $\mathsf{0}$ to (x, y, z) is geodesic means that $\pi(\alpha)$ is the shortest path in $\mathfrak{m}$ from $(0, 0)$ to (x, y) enclosing (nonnegative) balayage area z. By central symmetry of the isoperimetrix, we can solve the problem for minimal signed area by traversing the isoperimetrix clockwise rather than counterclockwise. Therefore the minimal signed

area enclosed by a path of length 1 from the origin to (x, y) is given by $-A(-x, -y) = -A(x, y)$, which tells us that the graph of $\mathbf{z}$ is symmetric over the xy plane.

Continuity of $A(x, y)$ on Q is is immediate from the continuity of $\mathsf{trace}(t, T)$ in its parameters. It only remains to show that the restriction of $A(x, y)$ to each Q_{ij} is a quadratic polynomial.

Observe that the problem of finding the trace path to a point $(x, y) \in Q_{ij}$ amounts to finding scalars s_1, s, s_2 (with $s_1, s_2 \leq s$) such that

$$(x, y) = s_1 \mathsf{w}_i + s(\mathsf{w}_{i+1} + \cdots + \mathsf{w}_{j-1}) + s_2 \mathsf{w}_j,$$

where w_i are the vectors defining the sides of I, as in Figure 2.



FIGURE 2. The balayage area of a path that follows along a scaled copy of the isoperimetrix is given by a quadratic expression in s_1, s_2, s.

The area enclosed is equal to that of a polygon of fixed shape with $j - i$ sides of length proportional to s, so that its area is proportional to s^2, plus two triangles of area proportional to $s_1 \cdot s$ and $s_2 \cdot s$, respectively. There are three linear equations relating s_1, s_2, s, given by the total arclength equaling one and the Euclidean displacement between endpoints equaling x and y in the horizontal and vertical direction, respectively. Thus we can solve to get expressions for s_1, s_2, s that are linear in x, y; this means that the area is quadratic in x, y, as required. □

By the definition of a CC metric, the sphere of radius t based at $\mathsf{0}$ is precisely $\delta_t \mathcal{S}$ (so $\mathcal{S}$ is homeomorphic to a 2-sphere), and the spheres with other centers are obtained by pushing these spheres around by left-multiplication in $H(\mathbb{R})$.

We want to break down the sphere, which is the graph of $\mathbf{z}$ over Q, into the graphs over the various quadrilaterals Q_{ij}. We will call these the *panels* of the sphere, and define them by

$$\mathsf{Panel}_{ij} := \{(x, y, z) : \quad (x, y) \in Q_{ij}, \quad z = \pm A(x, y)\}, \qquad j \neq i + 1,$$

and

$$\mathsf{Panel}_{i,i+1} := \{(x, y, z) : \quad (x, y) \in Q_{i,i+1}, \quad z \in [-A, A]\},$$

so that $\mathcal{S} = \bigcup_{1 \leq i,j \leq 2N} \mathsf{Panel}_{ij}$. Note that we can leave out the Panel_{ii} without changing this equality, because they are contained in the union of the other panels. The $\mathsf{Panel}_{i,i+1}$ project to edges of L, and the fact that $\mathbf{z}$ is quadratic over the interior implies that they are bounded above and below by parabolas in a plane through that edge that is

perpendicular to $\mathfrak{m}$. We will call these the *side panels* of $\mathcal{S}$ (these are the panels cut away in Figure 3).

The combinatorics of trace paths on quadrilaterals dictates the shape of CC geodesics in the following straightforward way. Let the *footprint map* $\mathsf{Foot} : H(\mathbb{R}) \to Q$ be defined by

$$\mathsf{Foot}(\mathsf{x}) := \pi \circ \delta_{1/d}(\mathsf{x}) = \delta_{1/d} \circ \pi(\mathsf{x}),$$

where $d = d_{\mathrm{CC}}(\mathsf{x}, \mathsf{0})$. Note that all geodesics from $\mathsf{0}$ to any point are just dilations of geodesics from $\mathsf{0}$ to $\mathcal{S}$. These in turn are lifts of trace paths and beelines, whose shape is determined by which quadrilateral contains the projection from $\mathcal{S}$ to Q. That is, the shapes of all possible geodesics from $\mathsf{0}$ to x are determined by which quadrilateral contains $\mathsf{Foot}(\mathsf{x})$.

Let the *regular points* in the unit sphere be the union of the panels over the quadrilaterals interior to Q and the *unstable points* be those in side panels.

$$\mathcal{S}_{\mathsf{reg}} = \bigcup_{j \neq i+1} \mathsf{Panel}_{ij} \quad ; \qquad \mathcal{S}_{\mathsf{uns}} = \bigcup \mathsf{Panel}_{i,i+1} \ ,$$

so that $\mathcal{S} = \mathcal{S}_{\mathsf{reg}} \cup \mathcal{S}_{\mathsf{uns}}$. Recall that the points of Q° are reached only by trace paths, while points of L are reached by families of beelines. We have

$$\Delta\, \mathcal{S}_{\mathsf{reg}} = \overline{\mathsf{Foot}^{-1}(Q^\circ)} \quad ; \qquad \Delta\, \mathcal{S}_{\mathsf{uns}} = \mathsf{Foot}^{-1}(L).$$

That is, the regular/unstable distinction divides the space $H(\mathbb{R})$ according to which of the two types of geodesic reaches each point.



FIGURE 3. Two views of the limit shape $\mathcal{S}$ for the hexagonal generators, with the quadrilateral decomposition of the footprint Q reproduced for reference. Only the regular part, $\mathcal{S}_{\mathsf{reg}}$, is shown. The unstable part, $\mathcal{S}_{\mathsf{uns}}$, is cut away in the picture: it is made up of six flat panels perpendicular to the xy-plane that are bounded by parabolas.

Let the *volume subtended by a panel* be the volume of the region obtained by coning off to the origin by dilation. We will see that the the decomposition of the sphere into regular and unstable points gives us useful invariants coming from volume:

$$V_{\mathsf{reg}} = \mathrm{vol}(\widehat{\mathcal{S}}_{\mathsf{reg}}) \quad ; \qquad V_{\mathsf{uns}} = \mathrm{vol}(\widehat{\mathcal{S}}_{\mathsf{uns}}).$$

Recalling that V is the volume of the unit ball, we have $V = V_{\mathsf{reg}} + V_{\mathsf{uns}}$.

Remark 11 (Rationality)**.** We note that Breuillard carried out a full description for the standard generators of the kind given in this section, and indicated key elements of such a description for general S, in [2, Prop 9.1]. For instance, it is stated without argument there that the balayage function should be piecewise quadratic. Breuillard sketches an argument that V (the volume of the unit ball for any limit metric) is rational, noting that Q has integer vertices, so its polar dual and therefore the isoperimetrix I must have rational vertices (since $Q^* = \{\mathsf{x} \in \mathfrak{m} : \ \mathsf{x}\cdot\mathsf{y} \le 1 \quad \forall \mathsf{y} \in Q\}$). Following this line in our language, we see that the vectors w_i have rational projections to each coordinate direction as well as rational length in the L-norm (though not in the Euclidean norm), and so the linear relations between x, y and s_1, s_2, s described in the proof of Theorem 10 are rational, which ensures that the balayage area over Q_{ij} is a rational quadratic polynomial in x and y. Furthermore, the vertices of the quadrilaterals Q_{ij} are rational (when a trace path begins and ends at a vertex, and has arclength 1, then each coordinate of the endpoint is a sum of rational numbers, scaled by a rational number). Thus the volume subtended by each panel is given by an integral over a region bounded entirely by graphs of rational quadratic polynomials (both the balayage area and the tracks of the dilation) over rational polygonal domains, so each such volume is in fact rational.

Remark 12 (Comparing metrics)**.** Notice the following fact that easily follows from Pansu's theorem from this geometric description of limit shapes. For an arbitrary generating set S of $H(\mathbb{Z})$, despite the fact that the word-length of generators is one by definition and the CC metric is the limit of the rescaled word metrics, it is easily possible for elements of S to be artibrarily far from $\mathsf{0}$ in CC distance: one can have $d_{\mathrm{cc}}(\mathsf{a}, \mathsf{0}) \gg 1 = |\mathsf{a}|_S$ for some $\mathsf{a} \in S$, meaning that the generator lies far outside of $\mathcal{S}$. This is because d_{cc} depends only on $\pi(S)$ and not on the z-coordinates of the generators. However, if $\pi(\mathsf{a})$ is not a vertex of L, we will eventually have $d_{\mathrm{cc}}(\mathsf{a}^n, \mathsf{0}) \le |\mathsf{a}^n|_S$ because the z-coordinate of a^n grows only linearly, while the height of $\delta_n\mathcal{S}$ over $\pi(\delta_n\mathsf{a})$ grows quadratically in n. It is only over the vertices of L that the height of $\mathcal{S}$ is zero. In that case, we see that $d_{\mathrm{cc}}(\mathsf{a}^n, \mathsf{0}) \le |\mathsf{a}^n|_S + 1$ for sufficiently large n, because the dilation causes the curves defining the sphere near its corners to get more and more vertical as the radius gets large. This establishes that there is a constant $K = K(S)$ such that

$$d_{\mathrm{cc}}(\mathsf{x}, \mathsf{0}) \ \le \ |\mathsf{x}|_S + K$$

for all $\mathsf{x} \in H(\mathbb{Z})$, which is half of Krat's bounded difference theorem (Thm 3).

3.2. **Nonuniqueness and regular points.** The union of the degenerate quadrilaterals has a geometric significance. Recall that the quadrilateral Q_{ii} is a line segment based at the origin along with a vertex. Let v_i' denote the other endpoint of the line segment emanating from $\mathsf{0}$. The other degenerate quadrilaterals cover the boundary L, whose vertices are called v_i. Then, with the convention that $Q_{2N,2N+1} = Q_{2N,1}$, let

$$\mathsf{N}_0 := \bigcup_{1\le i\le 2N} Q_{ii} \setminus \{\mathsf{v}_i'\} \quad ; \qquad L_0 := \bigcup_{1\le i\le 2N} Q_{i,i+1} \setminus \{\mathsf{v}_i\} = L \setminus \{\mathsf{v}_i\}.$$

Now define $\mathsf{N} := \mathsf{N}_0 \cup L_0$.

Proposition 13 (Uniqueness of CC geodesics)**.** *A point $\mathsf{x} \in \mathcal{S}$ is reached by more than one geodesic of length one based at the origin if and only if $\pi(\mathsf{x}) \in \mathsf{N}$.*

Thus a point $\mathsf{x} \in H(\mathbb{R})$ is reached by more than one geodesic based at the origin if and only if $\mathsf{Foot}(\mathsf{x}) \in \mathsf{N}$.

Proof. First, we consider the case of trace paths. It is impossible for a convex polygon (in this case λI_1) to have two different interior chords which are parallel, have the same length, and subtend the same perimeter on each side. Thus the only way this non-rigidity can occur is for the chord α to be contained in a side of λI_1. But this is precisely the case when the start and end points of trace fall on the same side, which was examined in the discussion of Q_{ii}. The vertices of L have trace paths enclosing zero area, so the geodesics are unique in that case. Finally, if $\pi(\mathsf{x})$ is interior to a side of L, then geodesics are lifts of beelines that are not straight lines. These can always be perturbed to nearby curves with the same endpoints, length, and enclosed area. □

From now on we call N the *nonuniqueness locus* in Q.

Theorem 14 (Probability of unique geodesics)**.** *In $H(\mathbb{R})$ with the* CC *metric induced by a polygon L, the asymptotic density of unique geodesics equals V_{reg}/V.*

That is, choose x uniformly in the ball of radius r and consider the probability that there is only one geodesic from $\mathsf{0}$ to x. For every r, this probability equals V_{reg}/V.

Proof. The only instances of nonuniqueness occur over N, as we have seen. But $\mathsf{Foot}^{-1}(\mathsf{N}_0)$ has no volume, and none of the volume of $\Delta\,\mathcal{S}_{\mathsf{uns}}$ is contributed by the vertices. Thus the probability of being reached by distinct geodesics is precisely the proportion of the volume of the unit ball that is in the unstable part (i.e., subtended by the side panels). □

Example 15 (Volume calculations in ℓ^1 norm)**.** Let $\|\cdot\|_L$ be the ℓ^1 norm on $\mathfrak{m}$ and give $H(\mathbb{R})$ the corresponding CC metric. We find that the volume subtended by panels with 4-sided combinatorics is 13/216, the volume subtended by 3-sided panels is 11/54, and the volume subtended by the side panels is 1/6. Thus $V_{\mathsf{reg}} = 13/216+11/54 = 19/72$ and $V_{\mathsf{uns}} = 1/6$. This adds up to give the total volume of the unit ball as $V = 31/72$, which agrees with calculations by Breuillard [2] and Stoll [11] of the volume growth.

Thus for this standard polygonal CC metric, the density of points reached by nonunique geodesics is 12/31, or about 39%, versus $19/31 \approx 61\%$ for unique geodesics.

4. Geodesics in the word metric

4.1. **Comparison of word and CC geodesics.** Recall that the *Hausdorff distance* $d_{\mathsf{Haus}}(\alpha,\beta)$ between a pair of sets is the smallest $\epsilon \geq 0$ such that each set lies in the ϵ-neighborhood of the other. Below, we will talk about paths and the images of those paths interchangeably, to make sense of the Hausdorff distance between paths. The goal of this section is make precise the following statement: "CC geodesics approximate word geodesics."

Lemma 16 (Continuity of CC geodesics)**.** *Fix an arbitrary $\rho > 0$. Suppose we are given $\mathsf{x} \in \mathcal{S}$ such that $\pi(\mathsf{x}) \notin \mathcal{N}_\rho(\mathsf{N}_0)$ and a* CC *geodesic α from $\mathsf{0}$ to x. Then for every $\epsilon > 0$ there exists $\delta > 0$ such that whenever $\mathsf{y} \in \mathcal{S}$ with $d_{\mathrm{CC}}(\mathsf{x},\mathsf{y}) < \delta$, there exists a* CC *geodesic β from $\mathsf{0}$ to y with $d_{\mathsf{Haus}}(\alpha,\beta) < \epsilon$.*

Proof. This is clear for points interior to any regular panel. For points x in the boundaries of regular panels, one can see from the combinatorial description that a sequence of points approaching x from any panel must have geodesics approaching the unique geodesic to x (see Figure 1). Suppose x is in the interior of a side panel and y is close to x. If $\pi(\mathsf{x}) = \pi(\mathsf{y})$, then homotope $\pi(\alpha)$ to a path $\overline{\beta}$ with the same endpoints such that the tracks of individual points are small. As long as every tangent vector of every point

on every intermediate path points towards the same side, this is a homotopy through beelines. If $\pi(\mathsf{x}) \neq \pi(\mathsf{y})$, then we can concatenate this homotopy with another similar homotopy ending at a path from the origin to $\pi(\mathsf{y})$ without changing area much. Either way, lift $\overline{\beta}$ to a geodesic β ending at y which is close to α. □

This continuity statement is false if $\pi(\mathsf{x}) \in \mathsf{N}_0$. For instance, take x to be the point on $\mathcal{S}$ above $\mathsf{0}$ and $\{\mathsf{y}_n\}$ to be a sequence of points interior to a panel, converging to x. Then for each n, there is a unique geodesic γ_n from $\mathsf{0}$ to y_n; these converge to a particular one of the many geodesics from $\mathsf{0}$ to x, which include the lifts of $\mathsf{trace}_{t,t+1}$ for all $t \in [0,1]$. These other geodesics to x are thus not closely approximated by geodesics to the y_n even as $\mathsf{y}_n \to \mathsf{x}$.

Let S be a finite generating set for $H(\mathbb{Z})$, and K be the maximum of $d_{\text{cc}}(\mathsf{0}, \mathsf{a})$ over all generators $\mathsf{a} \in S$. Minimal-length spellings $\mathsf{w} = \mathsf{a}_1 \cdots \mathsf{a}_n$ written in letters from S may be realized as *discrete geodesics* $\mathsf{x}_0, \mathsf{x}_1, ..., \mathsf{x}_n$ in $H(\mathbb{R})$ where $\mathsf{x}_0 = \mathsf{0}$, $\mathsf{x}_i = \mathsf{a}_1 \cdots \mathsf{a}_i$, and the CC distance between successive points is no more than K. We will say that the discrete geodesic $\mathsf{x}_0, \mathsf{x}_1, ..., \mathsf{x}_n$ is *ϵ-linearly tracked* (or just ϵ-tracked) by an admissible path α if they have the same endpoints and the set $\{\mathsf{x}_i\}$ stays inside the ϵn-neighborhood of α.

Lemma 17 (Geodesic spellings vs. admissible paths)**.** *For any $\epsilon > 0$ and $\mathsf{x} \in H(\mathbb{Z})$ with sufficiently large $n = |x|$, every discrete geodesic from $\mathsf{0}$ to x is ϵ-linearly tracked by an admissible path of length $n + O(\sqrt{n})$.*

Proof. Let $z(\mathsf{x})$ denote the height (i.e., the z-coordinate) of $\mathsf{x} \in H(\mathbb{R})$, and $M := \max z(S)$. Let $\mathsf{x}_0, ..., \mathsf{x}_n$ be a discrete geodesic from $\mathsf{0}$ to $\mathsf{x} \in H(\mathbb{R})$ in letters from S. Define a polygonal path γ in $\mathfrak{m}$ by connecting vertices $\pi(\mathsf{x}_0), \ldots, \pi(\mathsf{x}_n)$ with straight lines.

The curve γ has a unique admissible lift $\overline{\gamma}$; its endpoints are at $\mathsf{0}$ and at a point $\mathsf{y} \in H(\mathbb{R})$ which lies in the same vertical line as x. Since $\pi(S) \subset Q$, the curve γ lies in nQ, and since it is made up of straight segments of L-length ≤ 1, the length of γ (and hence $\overline{\gamma}$) is bounded above by n. We will establish the following equation:

$$z(\mathsf{x}) = z(\mathsf{y}) + \sum_{i=1}^{n} z(\mathsf{a}_i) \tag{†}$$

This says that the height of x is the balayage area of γ plus the heights of all of the letters used in the spelling.

Equation (†) can be proven by induction. The group law, in exponential coordinates, says

$$(x, y, z)(x', y', z') = \left(x + x',\ y + y',\ z + z' + \frac{xy' - yx'}{2}\right).$$

In particular, if $\mathsf{x}_{n-1} = (x, y, z)$ and $\mathsf{a}_n = (x', y', z')$, then we see that the change in height between x_{n-1} and $\mathsf{x}_n = \mathsf{x}_{n-1}\mathsf{a}_n$ is equal to the height of a_n plus the area of an appropriate triangle in $\mathfrak{m}$. That means that the height of the spelling path differs from the height predicted by balayage area by precisely the height of the generators, as claimed in (†).

Now set $C = d_{\text{cc}}((0, 0, M), \mathsf{0})$, so that

$$d_{\text{cc}}(\mathsf{0}, (0, 0, Mn)) = d_{\text{cc}}(\mathsf{0}, \delta_{\sqrt{n}}(0, 0, M)) = C\sqrt{n}.$$

Let α be a path obtained by concatenating $\overline{\gamma}$ with a CC geodesic from y to x. From (†) it follows that

$$|z(\mathsf{x}_i) - z(\mathsf{y}_i)| \leq Mi \leq Mn,$$

for all i, where y_i is the point on $\overline{\gamma}$ corresponding to the vertex x_i. Therefore $d_{\text{CC}}(\mathsf{x}_i, \mathsf{y}_i) \leq C\sqrt{n}$, so the discrete geodesic lives in the $C\sqrt{n}$-neighborhood of α, which means that the path $Cn^{-1/2}$-tracks the spelling. Furthermore, the difference in lengths between α and $\overline{\gamma}$ is bounded by $C\sqrt{n}$. Thus the length of α is bounded above by $n + C\sqrt{n}$. On the other hand, Theorem 3 provided a global constant C' such $d_{\text{CC}}(\mathsf{0}, \mathsf{x}) \geq n - C'$. This proves that the length of α is $n + O(\sqrt{n})$. □

Lemma 18 (Tracking Lemma)**.** *For every $\epsilon > 0$, $\rho > 0$, and sufficiently large $\mathsf{x} \in H(\mathbb{Z})$, if $\mathsf{Foot}(\mathsf{x}) \notin \mathcal{N}_\rho(\mathsf{N}_0)$, then every discrete geodesic from $\mathsf{0}$ to x is ϵ-linearly tracked by a* CC *geodesic.*

Proof. Suppose otherwise; then there is an $\epsilon > 0$ such that for every $n \geq 0$, there exist points $\mathsf{x}_n \in H(\mathbb{Z})$ such that $\mathsf{Foot}(\mathsf{x}_n) \notin \mathcal{N}_\rho(\mathsf{N}_0)$, $|\mathsf{x}_n| \geq n$ and there are choices of discrete geodesics from $\mathsf{0}$ to x_n which stay $\epsilon|\mathsf{x}_n|$-far away from any CC geodesics to the same point. Let α_n be the admissible paths approximating these discrete geodesics, as described above.

Denote $\overline{\mathsf{x}}_n = \delta_{1/|\mathsf{x}_n|}\mathsf{x}_n$ and $\overline{\alpha}_n = \delta_{1/|\mathsf{x}_n|}\alpha_n$. The lengths of the $\overline{\alpha}_n$ are converging to 1 by Lemma 17. By passing to a subsequence, we may assume that the $\overline{\mathsf{x}}_n$ converge to a limit $\overline{\mathsf{x}} \in \mathcal{S}$ with $\pi(\overline{\mathsf{x}}) \notin \mathcal{N}_\rho(\mathsf{N}_0)$, and that the paths $\overline{\alpha}_n$ converge in Hausdorff distance to a geodesic $\overline{\alpha}$ from $\mathsf{0}$ to $\overline{\mathsf{x}}$ by Arzelà-Ascoli. By Lemma 16, there is a large enough n such that there is a geodesic $\overline{\beta}_n$ within a Hausdorff distance of $\epsilon/2$ from $\overline{\alpha}$, and for large enough n we know that $\overline{\alpha}_n$ is within $\epsilon/2$ of $\overline{\alpha}$. But then for $\beta_n = \delta_{|\mathsf{x}_n|}\overline{\beta}_n$, we have $d_{\mathsf{Haus}}(\alpha_n, \beta_n) < \epsilon|\mathsf{x}_n|$. So the discrete geodesics are ϵ-tracked by the CC geodesics β_n, giving us a contradiction. □

A key idea in above proof is that additive quasi-geodesics get close to true geodesics as the additive constant goes to zero. Again, this fails near N_0.



FIGURE 4. In these diagrams in $(\mathfrak{m}, \ell^1)$, we see a true geodesic α_n from $(0, 0)$ to (n, n) on the left and a path β_n with different combinatorics, enclosing the same area but with excess length, on the right. When these are rescaled so that their lifts terminate on the sphere of radius 1, the ratio of lengths goes to 1 while the Hausdorff distance is bounded away from zero.

Example 19. Let $\mathsf{x}_n = (n, n, \ n^4 - \frac{1}{2}n^2)$. There is a unique CC geodesic α_n from the origin to x_n, and its projection to the horizontal plane is shown in Figure 4 on the left. On the right is pictured a non-geodesic path β_n to x_n which has length $4n^2$ (while the true geodesic has length $L_n = 4n^2 - 2n$). Let $\mathsf{y}_n = \delta_{1/L_n}\mathsf{x}_n$ be the dilate of x_n that

lies in the unit sphere $\mathcal{S}$. Note that $\mathsf{y}_n \to (0, 0, 1/16)$ as $n \to \infty$. By scaling back the paths β_n by the same factor, we obtain paths whose lengths approach 1. However, the Hausdorff distance between α_n and β_n is $n^2 - n$, so after scaling they still have a Hausdorff distance of at least $1/4$. In the limit as $n \to \infty$, both kinds of paths limit to valid closed geodesics to $(0, 0, 1/16)$. But for any finite value of n, the path with the wrong combinatorics is far from any geodesic.

4.2. Instability of geodesics in $H(\mathbb{R})$ and $H(\mathbb{Z})$.

Definition 20. *For a geodesic space X with basepoint x_0, let $\mathcal{G}(x)$ be the set of all geodesics from x_0 to x. Then the* instability function $\mathbf{I}(x)$ *measures how far apart they can be, relative to the size of x:*

$$\mathbf{I}(x) := 2 \cdot \frac{\operatorname{diam} \mathcal{G}(x)}{d(x, x_0)} = \tfrac{2}{d(x,x_0)} \cdot \sup_{\alpha,\beta \in \mathcal{G}(x)} d_{\mathsf{Haus}}(\alpha, \beta),$$

where the metric on $\mathcal{G}(x)$ is given by Hausdorff distance.

We will say that a point x is ϵ-stable *if $\mathbf{I}(x) \le \epsilon$.*

In general, $\operatorname{diam} \mathcal{G}(x) \le d(x, x_0)/2$, because any point on a geodesic from x_0 to x is within $d(x, x_0)/2$ of one of the endpoints, and therefore within that distance of any other geodesic. Thus the factor of 2 in the definition normalizes $\mathbf{I}$ so that $0 \le \mathbf{I}(x) \le 1$ for any X, x_0, x.

Note that if X has unique geodesics, then $\mathbf{I} \equiv 0$. If X is δ-hyperbolic, then $\operatorname{diam} \mathcal{G}(x) \le \delta$ for all x, so $\mathbf{I}(x) \to 0$ as $x \to \infty$.

The asymptotic behavior of this instability function is far from being quasi-isometry invariant. Obviously $\mathbf{I} \equiv 0$ on the Euclidean plane $\mathbb{R}^2$, but the situation is quite different for $\mathbb{R}^2$ with a polygonal norm. For instance, in the ℓ^1 norm, the elements along the x and y axes have $\mathbf{I} = 0$, but the diagonal has $\mathbf{I}(a, a) = 1$. It is easily verified that with any generating set, the limit shape L divides the plane into sectors, and the instability is 0 on the boundaries of the sectors and 1 on the midlines, varying linearly between these extremes along the sides of L. In particular one can recover the vertex directions of L from the function $\mathbf{I}$. Since the word metric on $(\mathbb{Z}^2, S)$ is asymptotic to $\mathfrak{m}_L = (\mathbb{R}^2, \|\cdot\|_L)$, it is also true for word metrics that $\mathbf{I}$ detects the directions of significant generators.

The goal of this section is to study instability in the Heisenberg group. As one would expect, $H(\mathbb{Z})$ is more geodesically stable than $\mathbb{Z}^2$ but less stable than a hyperbolic group. More intriguingly, the stability depends on the generating set in a different way than for those other groups.

We will call an element $H(\mathbb{Z})$ regular if it is regular as an element of $H(\mathbb{R})$.

Lemma 21 (Stability of regular points)**.** *For every $\epsilon > 0$ and $\rho > 0$, all sufficiently large regular points $\mathsf{x} \in H(\mathbb{Z})$ with $\mathsf{Foot}(\mathsf{x}) \notin \mathcal{N}_\rho(\mathsf{N}_0)$ satisfy*

$$d_{\mathsf{Haus}}(g, h) < \epsilon |\mathsf{x}|$$

for all word geodesics g, h from $\mathsf{0}$ to x.

Proof. Let S be a finite generating set for $H(\mathbb{Z})$ and give $H(\mathbb{R})$ the corresponding CC metric. Let $T \ge 0$ be large enough so that when $|\mathsf{x}| \ge T$ and $\mathsf{Foot}(\mathsf{x}) \notin \mathcal{N}_\rho(\mathsf{N}_0)$ and g is a word geodesic from $\mathsf{0}$ to x, then there is a CC geodesic γ from $\mathsf{0}$ to x such that $d_{\mathsf{Haus}}(g, \gamma) < \frac{\epsilon}{2}|g| = \frac{\epsilon}{2}|\mathsf{x}|$. Since regular points have unique CC geodesics, we are done. □

Then since the proportion of $\Delta\mathcal{S}_{\mathsf{reg}}$ lying over $\mathcal{N}_\rho(\mathsf{N}_0)$ goes to zero as $\rho \to 0$, we can conclude from the point of view of geometric probability that all of the regular part of $H(\mathbb{R})$ is ϵ-linearly stable for every ϵ, so that Hausdorff distance between word geodesics is bounded above by *every* linear function of word length. That is, for regular points of $H(\mathbb{Z})$, word geodesics satisfy a sublinear fellow traveling property. This is much stronger control than one has in $\mathbb{Z}^d$, where only words that are powers of single generators can be ϵ-stable for every ϵ, so almost all points are unstable. At the other extreme, all elements sufficiently far from the basepoint in a hyperbolic group G are stable. The Heisenberg group is intermediate between these in terms of geodesic stability: only the regular part is ϵ-stable for all ϵ, and its measure depends nontrivially on the generators.

To sum up this comparison:

Theorem 22 (Instability Theorem)**.** *With respect to any finite generating set,*

$$\lim_{\epsilon\to 0}\mathrm{Prob}\big(\mathbf{I}(\mathsf{x}) \le \epsilon\big) = \begin{cases} 0, & X = \mathbb{Z}^d, \\ V_{\mathsf{reg}}/V, & X = H(\mathbb{Z}), \\ 1, & X \text{ unbounded, } \delta\text{-hyperbolic.} \end{cases}$$

5. Lattice point counting

Because the balls in the CC metric get "fat" (i.e., the volume of a 1-neighborhood of the sphere is lower-order than the volume of the ball), it is easy to see that the number of lattice points in a ball of large radius equals the volume of the ball (to first order). What is much more delicate is to count lattice points in an annular region between a sphere of radius n and a sphere of radius $n-1$.

As before, the footprint of the limit shape, Q, is covered by finitely many quadrilaterals Q_{ij}. For a subset U of $H(\mathbb{R})$, let $\Delta_n U = \{\delta_t(U) : n-1 < t \le n\}$ be the "annular" region between the $(n-1)$st and nth dilate of U.

Recall the functions $\mathbf{z}$ and A described in Theorem 10. These functions are related by $A = \max \mathbf{z}$; that is, A is the height (maximum z-coordinate) of the unit sphere in the CC metric over points in Q, or equivalently, it is the maximum balayage area of curves of L-length 1 from the origin to (x, y). We can compute the height of the CC sphere of radius n above $(x, y) \in nQ$ using the rule $H_n(x, y) = n^2 \cdot A(x/n, y/n)$. (Note that for fixed (x, y) and large enough n, this is continuous and strictly increasing in n.) The unit ball is compact, so A takes a maximum value A_{max} on Q (the largest height achieved by the unit sphere). Then for fixed (x, y), the height function $H_n(x, y)$ is bounded above by $A_{\mathsf{max}} \cdot n^2$. But this is not by itself good enough to get control on the height difference between the n-sphere and the $(n-1)$-sphere over a point, $h_n := H_n - H_{n-1}$.

Lemma 23 (Control on height)**.** *Take a polygonal* CC *metric on $H(\mathbb{R})$ and let $h_n(x, y)$ be the height difference between the spheres of radius n and $(n-1)$ over (x, y) for $(x, y) \in (n-1)Q$. Fix any nondegenerate quadrilateral Q_{ij} and let $\Omega_n \subset nQ_{ij}$ denote the complement of the 1-neighborhood of the boundary. Then $h_n = O(n)$ over Ω_n.*

Proof. First of all, for $\mathsf{x} \in \Omega_n$, one easily verifies that both $\frac{1}{n}\mathsf{x}$ and $\frac{1}{n-1}\mathsf{x}$ lie in Q_{ij}. Recall from the structure theorem (Thm 10) that the nonnegative part of $\mathcal{S}$ over Q_{ij} is the graph of a single quadratic polynomial in x, y ; denote this polynomial by $f(x, y) = a_1x^2 + a_2xy + a_3y^2 + b_1x + b_2y + c$. For all $(x, y) \in \Omega_n$, we have

$$\big(x, y,\ n^2 \cdot f(\tfrac{x}{n}, \tfrac{y}{n})\big) \quad \in \delta_n(\mathsf{Panel}_{ij})$$

so

$$h_n(x,y) = n^2 \cdot f(x/n, y/n) - (n-1)^2 \cdot f(x/n-1, y/n-1) = b_1 x + b_2 y + 2cn - c.$$

Since $(x,y) \in \Omega_n \subset nQ_{ij} \subset nQ$, both x and y are $O(n)$, and therefore h_n is as well. □

Theorem 24 (Counting Theorem)**.** *Consider the* CC *metric induced on* $H(\mathbb{R})$ *by a norm* $\|\cdot\|_L$ *on* $\mathfrak{m}$*, with* L *a polygon. Let* σ *be an arbitrary measurable subset of* $\mathcal{S}$*, the* CC *unit sphere. Then*

$$\#\left(H(\mathbb{Z}) \cap \Delta_n \sigma\right) = 4n^3 \operatorname{vol}(\hat{\sigma}) + O(n^2).$$

Proof. First we consider $\sigma \subset \mathcal{S}_{\mathsf{reg}}$. For this case, it suffices to treat small subsets σ which project to squares in $\mathfrak{m}$ whose closures are contained in the interior of a single quadrilateral:

$$U := \pi(\sigma) = (u, u+\epsilon) \times (v, v+\epsilon)$$

such that $[u, u+\epsilon] \times [v, v+\epsilon] \subset Q_{ij}^{\circ}$. Since the dilation δ_t is just a homothety on $\mathfrak{m}$, the dilate tU is a square for every t. The condition on the closure of U guarantees that when n is large enough, nU does not intersect the 1-neighborhood of the boundary of nQ_{ij}.

Consider the rectangle $U_n := nU \cap (n-1)U$, let $A_n := \Delta_n \sigma \cap \pi^{-1}(U_n)$ be the part of the nth annular shell that is vertically over U_n, and let $B_n := \Delta_n \sigma \setminus A_n$ be the rest (see Figure 5).



FIGURE 5. This shows an xz-plane slice, with U_n along the x-axis, A_n in blue, and B_n in red. We have $A_n \cup B_n = \Delta_n \sigma$; we want to count the lattice points in that entire region. The A_n lattice points are easier to count, and the number in B_n is lower-order.

We will make the following chain of comparisons:

$$\#\left(H(\mathbb{Z}) \cap \Delta_n \sigma\right) \sim \#\left(H(\mathbb{Z}) \cap A_n\right) \sim \operatorname{vol}(A_n) \sim \operatorname{vol}(\Delta_n \sigma) \sim 4n^3 \operatorname{vol}(\hat{\sigma}),$$

while keeping track of the error term.

First we will estimate the number of lattice points in B_n, and its volume. The projection $\pi(B_n)$ is contained in a square annulus in $\mathfrak{m}$ of width $O(1)$ and sidelength $O(n)$. Thus the projection contains at most $O(n)$ lattice points (x,y). By the previous

lemma, the difference in heights between $\delta_n\mathcal{S}$ and $\delta_{n-1}\mathcal{S}$ over each of these points is at most $O(n)$, so the total number of lattice points in B_n is at most $O(n^2)$. Thus

$$\#(H(\mathbb{Z}) \cap \Delta_n\sigma) = \#(H(\mathbb{Z}) \cap A_n) + O(n^2).$$

Likewise, the volume of B_n is bounded above by integrating $h_n(x,y)$ over $(x,y) \in \pi(B_n)$, so $\mathrm{vol}(B_n) = O(n^2)$. This shows that $\mathrm{vol}(A_n) = \mathrm{vol}(\Delta_n\sigma) + O(n^2)$. For the comparison on the far right, we have

$$\mathrm{vol}(\Delta_n\sigma) = \mathrm{vol}(\delta_n\hat{\sigma}) - \mathrm{vol}(\delta_{n-1}\hat{\sigma}) = \left[4n^3 + O(n^2)\right]\mathrm{vol}(\hat{\sigma}),$$

because δ_t expands volume by t^4.

All that remains is to compare $\#(H(\mathbb{Z}) \cap A_n)$ with $\mathrm{vol}(A_n)$. Fix attention on an integer point $(x,y) \in U_n$. For all the lattice points in the vertical line over (x,y), let us count the number in $\Delta_n\sigma$. We find $\#(H(\mathbb{Z}) \cap A_n) = (\sum_{U_n \cap \mathbb{Z}^2} h_n) + O(n^2)$, because $\#(U_n \cap \mathbb{Z}^2) = O(n^2)$ and there is bounded error on each line. On the other hand, $\mathrm{vol}(A_n) = \int_{U_n} h_n$. Regard the sum as a Riemann sum approximation to the integral over the integer squares. Since $h(n)$ is linear, the Riemann sum gives an equal error on each of the $O(n^2)$ squares, for total error of order n^2.

Next we deal with the case $\sigma \subset \mathcal{S}_{\mathsf{uns}}$, where it suffices to consider $\sigma = \gamma \times [h, h+\epsilon]$ with γ a subinterval of an edge of L. Let $\Sigma = \Delta\gamma$ denote the sector of $\mathfrak{m}$ determined by γ, and let U_n be the trapezoid in Σ between $n\gamma$ and $(n-1)\gamma$. We see that $\Delta\sigma = \bigcup_{t\geq 0} t\gamma \times [ht^2, (h+\epsilon)t^2]$. Let the height difference from top to bottom at (x,y) be denoted $T(x,y) = \epsilon \cdot \|(x,y)\|_L$.

We shall make the comparison

$$\#(H(\mathbb{Z}) \cap \Delta_n\sigma) \sim \mathrm{vol}(\Delta_n\sigma)$$

by considering the Riemann sum approximation of $\int_{U_n} T$ by $\sum_{U_n \cap \mathbb{Z}^2} T$. For any $\mathsf{u}, \mathsf{v} \in U_n$, the L-norm is between $n-1$ and n, so $T(\mathsf{v}) - T(\mathsf{u}) = \epsilon\|\mathsf{v}\|_L - \epsilon\|\mathsf{u}\|_L \leq 2\epsilon n$, which gives a bound on the error per term in the Riemann sum. Since U_n has bounded width, its area and $\#\mathbb{Z}^2 \cap U_n$ are both $O(n)$. Putting these factors together, we get that the total error is $O(n^2)$, so we have shown that $\#(H(\mathbb{Z}) \cap \Delta_n\sigma) = 4n^3\,\mathrm{vol}(\hat{\sigma}) + O(n^2)$, as in the regular case. □

6. Limit measure in the standard generators

To get more precise results that allow averaging over spheres and not just balls, we focus attention on $H(\mathbb{Z})$ with the standard generating set $S = \mathsf{std} = \pm\{\mathsf{e}_1, \mathsf{e}_2\}$ and consider the word metric with respect to those generators. The goal for this section is to show for this special case that counting measure on discrete spheres in the word metric limits to cone measure (with respect to Heisenberg dilation) on the limit shape $\mathcal{S}$.

Theorem 25 (Limit measure for standard generators)**.** *Consider the* CC *metric induced by the ℓ^1 norm on $\mathfrak{m}$, which is the limit metric for $(H(\mathbb{Z}), \mathsf{std})$. For any measurable set $\sigma \subset \mathcal{S}$,*

$$\#(S_n \cap \Delta\sigma) = n^3\,\mathrm{vol}(\hat{\sigma}) + O(n^2),$$

and therefore

$$\lim_{n\to\infty} \frac{\#(S_n \cap \Delta\sigma)}{\#S_n} = \frac{\mathrm{vol}(\hat{\sigma})}{\mathrm{vol}(\hat{\mathcal{S}})}.$$

Thus for the discrete spheres S_n in $(H(\mathbb{Z}), \mathsf{std})$, the counting measure on $\delta_{1/n}(S_n)$ converges to cone measure on the square $L \subset \mathfrak{m}$.

To prove this, we will study the distribution of points in S_n, finding that the set S_n has points on the line $\{(x,y,t) : t \geq 0\}$ if and only if x, y are integers and $x+y+n$ is even; in that case it contains all lattice points between $\delta_n(\mathcal{S})$ and $\delta_{n-2}(\mathcal{S})$. This is illustrated in Figure 6.



FIGURE 6. This figure depicts a cross-section of $H(\mathbb{R})$, with the points of the sphere S_{16} in $(H(\mathbb{Z}), \mathsf{std})$ marked and three dilates of the CC sphere shown for comparison. As will be proven in this section, S_n has very nearly as many points as the lattice points contained in $\Delta_n\mathcal{S}$ (the shaded region): it has twice the points on half the lines.

Lemma 26 (CC distance versus height)**.** *For any polygonal* CC *metric, distance from the origin is an increasing function of height: regarded as a function of $t \geq 0$, the value $d_{\text{CC}}\big((x,y,t),\mathsf{0}\big)$ is continuous, stays constant on an interval $[0, T(x,y)]$, and is strictly increasing thereafter.*

Proof. Let $\mathsf{x}_t = (x,y,t)$. The distance of a point x from $\mathsf{0}$ is the value d such $\delta_{1/d}(\mathsf{x}) \in \mathcal{S}$. If $\mathsf{x}_t \in \Delta\,\mathcal{S}_{\mathsf{uns}}$, then the dilate of x_t that hits $\mathcal{S}$ will intersect a side panel, so the distance of $\mathsf{x}_t = (x,y,t)$ to the origin depends only on x, y. Now $\Delta\,\mathcal{S}_{\mathsf{uns}}$ is a closed set with interior which contains $(x,y,0)$ for every $(x,y) \in \mathfrak{m}$. Thus $(x,y,t) \in \Delta\,\mathcal{S}_{\mathsf{uns}}$ for a closed interval of times $[0, T(x,y)]$.

Now consider x_t and x_s with $s > t$. These have the same projection to $\mathfrak{m}$, and so do $\delta_r(\mathsf{x}_t)$ and $\delta_r(\mathsf{x}_s)$ for any r. Thus if $\delta_{1/d}(\mathsf{x}_t) \in \mathcal{S}$, we must have $\delta_{1/d}(\mathsf{x}_s)$ vertically above $\mathcal{S}$, meaning that $d_{\text{CC}}(\mathsf{x}_s, \mathsf{0}) > d$. □

Note also that if α is a path in a Cayley graph for $H(\mathbb{Z})$ (i.e., a word written in terms of the generators), then there is a corresponding admissible path in $H(\mathbb{R})$ which ends at the same point and whose CC length is the same as the word-length in that word metric. (See the proof of Lemma 17 for the construction, noting that the extra height created by generators is zero for $S = \mathsf{std}$, or whenever the generators lie in $\mathfrak{m}$.) That is, word paths are realizable as admissible paths in an obvious way, as in Figure 7.

FIGURE 7. Comparing geodesics: $\mathsf{x} = (4, 0, 3)$ is reached by a three-sided CC geodesic of length $5\frac{1}{2}$, while its word-length in the standard generators is 6.

However, the CC geodesic in $H(\mathbb{R})$ between two points in $H(\mathbb{Z})$ is often shorter than the geodesic in the word metric, as seen in the figure: the CC geodesics need not have corners at integer points, even if they begin and end at integer points.

Lemma 27 (Word length versus height)**.** *For any $(x, y) \in \mathbb{Z}^2$, let $\epsilon = \epsilon(x, y) = 1/2$ if x, y odd and $\epsilon = 0$ otherwise, so that $(x, y, \epsilon + m) \in H(\mathbb{Z})$ for all $m \in \mathbb{Z}$. Then $|(x, y, \epsilon + m)|_{\mathsf{std}}$ is nondecreasing for $m = 0, 1, 2, \ldots$.*

Proof. Suppose without loss of generality that (x, y) is in the first quadrant of $\mathfrak{m}$. Let g be a geodesic in $(H(\mathbb{Z}), \mathsf{std})$ from the origin to a point $\mathsf{x} = (x, y, z)$ where $z > 0$. Then z is equal to the signed area enclosed between g and the straight chord from $\mathsf{0}$ to $\pi(\mathsf{x})$. The path contains only e_1 moves and e_2 moves, and there must be some subword $\mathsf{e}_1\mathsf{e}_2$ or else the signed area would be nonpositive. But then let g' be the same path with some $\mathsf{e}_1\mathsf{e}_2$ subword replaced by $\mathsf{e}_2\mathsf{e}_1$. This has the same length and its balayage area is one less, so it is a path in $H(\mathbb{Z})$ from $\mathsf{0}$ to $(x, y, z-1)$. This may no longer be geodesic, but its length provides an upper bound: $|(x, y, z-1)| \leq |(x, y, z)|$. □

Next, it is quite easy to see that there is a parity condition on lengths of paths from the origin to (x, y, z).

Lemma 28 (Parity)**.** *For any path of length n in $H(\mathbb{Z})$ from 0 to (x, y, z), n has the same parity as $x + y$.*

Proof. All paths in $(H(\mathbb{Z}), \mathsf{std})$ are of the form $\Pi \mathsf{e}_1^{a_i} \mathsf{e}_2^{b_i}$, with $\sum a_i = x$, $\sum b_i = y$, and length $n = \sum |a_i| + |b_i|$. Since $r + |r|$ is even for all integers, it follows that $x + y + n \in 2\mathbb{Z}$.

Geometrically, this is just the observation that if a path in $\mathfrak{m}$ to (x, y) goes horizontally past x or vertically past y, it will have to backtrack by the same number of steps. □

For fixed $(x, y) \in \mathbb{Z}^2$ and $n \in \mathbb{N}$, let us count the points $(x, y, z) \in S_n$. We first note that $\pi(\delta_n \mathcal{S}) = nQ$, so there are no points of S_n over (x, y) if $\|(x, y)\|_L > n$.

Theorem 29 (Word length in terms of CC)**.** *Fix $(x, y) \in \mathbb{Z}^2$. For $z \geq 0$ in $\mathbb{Z} + \epsilon$, let n be the unique integer with the same parity as $x + y$ such that $n - 2 < d_{\mathrm{cc}}((x, y, z), \mathsf{0}) \leq n$. Then*

$$|(x, y, z)|_{\mathsf{std}} = n.$$

Stated another way, this formula says that for all $\mathsf{x} \in H(\mathbb{Z})$,

$$d_{\mathrm{cc}}(\mathsf{x}, \mathsf{0}) \leq |\mathsf{x}|_{\mathsf{std}} < d_{\mathrm{cc}}(\mathsf{x}, \mathsf{0}) + 2,$$

and describes which of the two integers in that range is the correct value.

Proof. We will restrict ourselves to the half-space $z \geq 0$ without loss of generality; the situation on the other half-space is obtained by reflection. The point $(x, y, \epsilon + m)$ lies in the unstable part of $H(\mathbb{R})$ for $0 \leq m \leq T$, then in a sector with three-sided combinatorics

for some $T \le m \le T'$, and finally in a sector with four-sided combinatorics for $m \ge T'$. (See Figure 1.) We consider those three cases separately.

Fix $n \ge 0$ of the same parity as $x+y$. By Lemma 26, there is a unique real value $r \ge 0$ such that (x,y,r) is reached by a CC geodesic of length n. Take that r and suppose that the geodesic γ is three-sided. Note that $\pi(\gamma)$ is a path in $\mathfrak{m}$ moving only horizontally and vertically; without loss of generality, γ has the form $\mathsf{e}_2^{-t}\mathsf{e}_1^{x}\mathsf{e}_2^{y+t}$, where $t>0$ and $x+2t+y=n$. But we know $x+y$ has the same parity as n, so it follows that t is an integer. Thus γ is the realization of a word geodesic, so $|(x,y,r)| = d_{\mathrm{cc}}((x,y,r),0)$.

Now assume that γ is a 4-sided geodesic from the origin to (x,y,r), with length n. Without loss of generality, γ has the form $\mathsf{e}_2^{y-x-t}\mathsf{e}_1^{x+t}\mathsf{e}_2^{x+t}\mathsf{e}_1^{-t}$, where $t \ge 0$ and $3x-y+4t=n$. Since $x+y+n$ is even, it follows that $4t$ is an integer with the same parity as $4x$, so t is either an integer or a half-integer. If t is an integer, then γ is the realization of a word geodesic and $|(x,y,r)| = d_{\mathrm{cc}}((x,y,r),0)$. If not, then let $s = t-1/2$ and $u = t+1/2$ be the nearest integers. We can form an integer path $\mathsf{e}_2^{y-x-u}\mathsf{e}_1^{x+s}\mathsf{e}_2^{x+u}\mathsf{e}_1^{-s}$ of length n by shortening the horizontal sides and lengthening the vertical sides by $1/2$. This encloses area $z = (x+s)(x+u) - \frac{1}{2}xy$, which is in $\epsilon+\mathbb{Z}$, so the endpoint (x,y,z) of the new path is in $H(\mathbb{Z})$. This area is smaller than $r = (x+t)^2 - \frac{1}{2}xy$ by $1/4$, so it must be the nearest lattice point. This new path may not be a geodesic, but it establishes the inequality $|(x,y,z)| \le n$. On the other hand, shortening all four sides of γ by $1/4$ produces a 4-sided CC geodesic to (x,y,r') of length $n-1$, and one easily checks that $r' < z < r$. This shows that $d_{\mathrm{cc}}((x,y,z),0) > n-1$, and it follows that $|(x,y,z)| = n$.

In these cases, we have established that the highest lattice point over (x,y) inside the closed CC ball of radius n has word-length exactly n. Now we want to show that the lowest lattice point over (x,y) outside the closed CC ball of radius $n-2$ also has word-length exactly n. But this point has CC norm larger than $n-2$, so it has word-length larger than $n-2$, and by parity considerations the word-length must be exactly n.

The last case is easy: for any unstable point $(x,y,z) \in H(\mathbb{R})$, its CC norm is $d_{\mathrm{cc}}((x,y,z),0) = x+y$. We claim that these points satisfy $|(x,y,z)|_{\mathsf{std}} = x+y$ as well. To see this, just note that starting with the two-sided path $\mathsf{e}_1^x\mathsf{e}_2^y$, which encloses area $\frac{1}{2}xy$ with length $x+y$, one can shave off area one unit at a time with generator swaps (replacing some occurrence of $\mathsf{e}_1\mathsf{e}_2$ with $\mathsf{e}_2\mathsf{e}_1$) until the area enclosed is z. This produces a word path whose length matches the CC geodesic, so it too is geodesic. □

Thus far, we have shown that if σ is a subset of $\mathcal{S}_{\mathsf{reg}}$, then $|S_n \cap \Delta\sigma| \sim |H(\mathbb{Z}) \cap \Delta_n\sigma|$, because S_n has twice as many points as $\Delta_n\sigma$, but on half of the lines (see Figure 6). On the other hand, $|S_n \cap \Delta\sigma| = |H(\mathbb{Z}) \cap \Delta_n\sigma|$ exactly for $\sigma \subset \mathcal{S}_{\mathsf{uns}}$. Putting these together and then using the Counting Theorem (Thm 24) to estimate the number of lattice points in $\Delta_n\sigma$, we obtain Theorem 25.

References

[1] S. Blachère, Word distance on the discrete Heisenberg group. Colloquium Mathematicum 95, vol. 1, 2003, 21–36.

[2] E. Breuillard, Geometry of groups of polynomial growth and shape of large balls. arXiv:0704.0095

[3] D.Yu. Burago, Periodic metrics. Representation theory and dynamical systems, 205–210, *Adv. Soviet Math.*, 9, Amer. Math. Soc., Providence, RI, 1992.

[4] H. Busemann, The isoperimetric problem in the Minkowski plane. AJM 69 (1947), 863–871.

[5] L. Capogna, D. Danielli, S. Pauls and J. Tyson, *An Introduction to the Heisenberg Group and to the Sub-Riemannian Isoperimetric Problem.* Birkhauser, Progress in Mathematics, 2007.

[6] P. Dani, The asymptotic density of finite-order elements in virtually nilpotent groups. Journal of Algebra 316 (2007) 54–78.
[7] M. Duchin, S. Lelièvre, and C. Mooney, The shape of spheres in free abelian groups, preprint.
[8] S.A. Krat, Asymptotic properties of the Heisenberg group. Journal of Mathematical Sciences, Vol. 110, No. 4 (2002) 2824–2840.
[9] P. Pansu, Croissance des boules et des géodésiques fermées dans les nilvariétés. Ergodic Theory Dynam. Systems 3 (1983), no. 3, 415–445.
[10] M. Shapiro, A geometric approach to the almost convexity and growth of some nilpotent groups. Math. Ann. 285, 601–624 (1989).
[11] M. Stoll, Rational and transcendental growth series for the higher Heisenberg groups. Invent. math. 126, 85109 (1996).
[12] M. Stoll, On the asymptotics of the growth of 2-step nilpotent groups. J. London Math. Soc. (2) 58 (1998) 38–48.

Department of Mathematics, 530 Church Street, Ann Arbor, MI 48109-1043
E-mail address: mduchin@umich.edu
URL: http://www.math.lsa.umich.edu/~mduchin/

Department of Mathematics, 530 Church Street, Ann Arbor, MI 48109-1043
E-mail address: cpmooney@umich.edu
URL: http://www.math.lsa.umich.edu/~cpmooney/